

Special Focus Report

Early gene changes induced by isotretinoin in the skin provide clues to its mechanism of action

Amanda M. Nelson, Wei Zhao, Kathryn L. Gilliland, Andrea L. Zaenglein, Wenlei Liu and Diane M. Thiboutot*

The Jake Gittlen Cancer Research Foundation and Department of Dermatology; The Pennsylvania State University College of Medicine; Hershey, PA USA

Key words: 13-*cis* retinoic acid, sebaceous gland, apoptosis, acne

Gene expression analysis of patient skin at one-week of oral isotretinoin therapy for severe acne illustrates the initial changes caused by 13-*cis* RA. Focus on the function of these significantly changed genes provides clues to its mechanism of action in skin. Significant gene changes included the upregulation of *lipocalin 2*, which encodes the protein neutrophil gelatinase associated lipocalin (NGAL). Detailed studies of NGAL's localization and function within the sebaceous gland demonstrated that NGAL mediates the apoptotic action of 13-*cis* RA on sebaceous glands, thereby suggesting that agents which selectively induce NGAL expression in sebaceous glands might represent therapeutic alternatives to the use of 13-*cis* RA for the treatment of acne.¹

In addition to NGAL, 13-*cis* RA affected the expression of numerous other genes in patient skin at one-week and in our sebocyte cell culture model, SEB-1. The results of these gene expression array analyses have been submitted to the National Center for Biotechnology Information Gene Expression Omnibus Database and are available under series accession number GSE10434. These early gene changes can be broadly categorized as tumor suppressors, protein processors and genes involved in transfer or binding of ions, amino acids, lipids or retinoids. Investigating the function of these potential candidate genes that mediate retinoid response in the skin provides a better understanding of the complete actions of 13-*cis* RA within skin with the goal of discovering safer alternatives to oral retinoid use in the treatment of acne and other skin diseases. For example, tazarotene induced gene 1 [*TIG1*, retinoic acid responder 1 (*RARRES1*)] encodes a tumor suppressor belonging to the latexin family of proteins, whose promoter is methylated (CpG island) and therefore silenced in a variety of cancers.²⁻⁵ The increase in expression of *TIG1* induced by 13-*cis* RA may mediate the known effects of this drug in chemoprevention of skin cancer in addition to the known suppressive effects on sebocyte proliferation.⁶⁻⁹ Furthermore, genes encoding both serine proteases and serine protease inhibitors were upregulated by 13-*cis* RA at one-week. Serine protease

inhibitors (serpins) are involved in tissue remodeling and control of inflammation.¹⁰ Increased expression of serpins have been reported in inflammatory processes such as psoriasis and inflammatory acne lesions.^{11,12} Since 13-*cis* RA is the most potent agent available to reduce severe inflammatory acne, it is possible that the upregulation of serpins, which in turn, scavenge pro-inflammatory proteins, mediates the anti-inflammatory effect of 13-*cis* RA. In support of this hypothesis, previous HPLC studies demonstrated that 13-*cis* RA can be isomerized to ATRA within sebocytes⁹ and it has been shown that SERPINA5 is capable of binding ATRA in vitro and may function in retinoid transport.^{13,14}

13-*cis* RA significantly increased expression of multiple members of the solute carrier family of proteins. Recent data have highlighted the importance of the solute carrier family of proteins in skin biology. For example, *SLC12A8*, which is upregulated within our one-week analysis, encodes for a sodium/potassium/chloride transporter that has recently been identified as a candidate gene for psoriasis susceptibility and is contained within the PSORS5 locus of CHR3q.^{15,16} Although, no mutations in *SLC12A8* have been linked to psoriasis, it is interesting to note that it is retinoid responsive. *ATP12A*, another significantly upregulated gene at one-week, encodes sodium/potassium ATPase, an integral membrane protein involved in solute transport responsible for the hydrolysis of ATP coupled with the exchange of hydrogen and potassium ions across membranes. This particular protein regulates ion flux into melanosomes and affects tyrosinase activity.¹⁷ Interestingly, another cation transporter (SLC24A5) has been identified as playing a prominent role in skin biology by regulating flux across the melanosome membrane in zebrafish and this gene links to skin color in humans.¹⁸ No other studies to date have examined the other solute carrier molecules and their relationship to normal human skin or skin disorders.

Retinoids are crucial to epidermal development and differentiation.¹⁹ S100 proteins modulate cellular differentiation, energy metabolism, cytoskeletal membrane interactions and cell cycle progression.²⁰ S100 protein family members are upregulated in our one-week patient array as well as our SEB-1 sebocytes after 13-*cis* RA treatment. Interestingly, the upregulated S100 proteins are induced by oxidative or inflammatory stress and *S100A7* (psoriasin) may function as a chemo-attractant agent for immune cells.²⁰ One can speculate that initial upregulation of S100 proteins by 13-*cis* RA is responsible for the "acne-flare" response observed in some patients receiving oral or topical retinoids for the treatment of their acne.

*Correspondence to: Diane M. Thiboutot; Department of Dermatology; HU14; The Pennsylvania State University College of Medicine; 500 University Drive; Hershey, PA 17033 USA; Tel.: 717.531.7437; Fax: 717.531.4821; Email: dthiboutot@psu.edu

Submitted: 01/15/09; Accepted: 02/09/09

Previously published online as a *Dermato-Endocrinology* E-publication:
<http://www.landesbioscience.com/journals/dermatoendocrinology/article/8107>

Approximately half of the significantly changed genes affected by 13-*cis* RA contained consensus sequences for RAR or RXR receptors. This study does not address the question of whether the changes in gene expression are the direct result of 13-*cis* RA, one of its metabolites or if retinoid receptors are involved. Instead, attention is focused on the function of these significantly changed genes and how these functions may provide clues to 13-*cis* RA within the sebaceous glands and whole skin.

References

1. Nelson AM, Zhao W, Gilliland KL, Zaenglein AL, Liu W, Thiboutot DM. Neutrophil gelatinase-associated lipocalin mediates 13-*cis* retinoic acid-induced apoptosis of human sebaceous gland cells. *J Clin Invest* 2008; 118:1468-78.
2. Mizuiri H, Yoshida K, Toge T, Oue N, Aung PP, Noguchi T, Yasui W. DNA methylation of genes linked to retinoid signaling in squamous cell carcinoma of the esophagus: DNA methylation of *CRBP1* and *TIG1* is associated with tumor stage. *Cancer Science* 2005; 96:571-7.
3. Zhang J, Liu L, Pfeifer GP. Methylation of the retinoid response gene *TIG1* in prostate cancer correlates with methylation of the retinoic acid receptor beta gene. *Oncogene* 2004; 23:2241-9.
4. Youssef EM, Chen XQ, Higuchi E, Kondo Y, Garcia-Manero G, Lotan R, et al. Hypermethylation and silencing of the putative tumor suppressor Tazarotene-induced gene 1 in human cancers. *Cancer Res* 2004; 64:2411-7.
5. Shutoh M, Oue N, Aung PP, Noguchi T, Kuraoka K, Nakayama H, et al. DNA methylation of genes linked with retinoid signaling in gastric carcinoma: Expression of the retinoid acid receptor β , cellular retinol-binding protein 1 and tazarotene-induced gene 1 genes is associated with DNA methylation. *Cancer* 2005; 104:1609-19.
6. Zouboulis CC, Korge B, Akamatsu H, Xia LQ, Schiller S, Gollnick H, et al. Effects of 13-*cis* retinoic acid, all-trans-retinoic acid and acitretin on the proliferation, lipid synthesis and keratin expression of cultured human sebocytes in vitro. *J Invest Dermatol* 1991; 96:792-7.
7. Zouboulis CC, Korge BP, Mischke D, Orfanos CE. Altered proliferation, synthetic activity and differentiation of cultured human sebocytes in the absence of vitamin A and their modulation by synthetic retinoids. *J Invest Dermatol* 1993; 101:628-33.
8. Zouboulis CC, Seltmann H, Neitzel H, Orfanos CE. Establishment and characterization of an immortalized human sebaceous gland cell line (SZ95). *J Invest Dermatol* 1999; 113:1011-20.
9. Nelson AM, Gilliland KL, Cong Z, Thiboutot DM. 13-*cis* Retinoic acid induces apoptosis and cell cycle arrest in human SEB-1 sebocytes. *J Invest Dermatol* 2006; 126:2178-89.
10. Silverman GA, Bird PI, Carrell RW, Church FC, Coughlin PB, Gettins PG, et al. The serpins are an expanding superfamily of structurally similar but functionally diverse proteins. Evolution, mechanism of inhibition, novel functions and a revised nomenclature. *J Biol Chem* 2001; 276:33293-6.
11. Takeda A, Higuchi D, Takahashi T, Ogo M, Baci P, Goetinck PF, et al. Overexpression of serpin squamous cell carcinoma antigens in psoriatic skin. *J Invest Dermatol* 2002; 118:147-54.
12. Trivedi NR, Gilliland KL, Zhao W, Liu W, Thiboutot DM. Gene array expression profiling in acne lesions reveals marked upregulation of genes involved in inflammation and matrix remodeling. *J Invest Dermatol* 2006; 126:1071-9.
13. Krebs M, Uhrin P, Vales A, Prendes-Garcia MJ, Wojta J, Geiger M, et al. Protein C inhibitor is expressed in keratinocytes of human skin. *J Invest Dermatol* 1999; 113:32-7.
14. Jerabek I, Zechmeister-Machhart M, Binder BR, Geiger M. Binding of retinoic acid by the inhibitory serpin protein C inhibitor. *Eur J Biochem* 2001; 268:5989-96.
15. Hewett D, Samuelsson L, Polding J, Enlund F, Smart D, Cantone K, et al. Identification of a psoriasis susceptibility candidate gene by linkage disequilibrium mapping with a localized single nucleotide polymorphism map. *Genomics* 2002; 79:305-14.
16. Hüffmeier U, Lascorz J, Traupe H, Böhm B, Schürmeier-Horst F, Ständer M, et al. Systematic linkage disequilibrium analysis of *SLC12A8* at PSORS5 confirms a role in susceptibility to psoriasis vulgaris. *J Invest Dermatol* 2005; 125:906-12.
17. Watabe H, Valencia JC, Yasumoto K, Kushimoto T, Ando H, Muller J, et al. Regulation of tyrosinase processing and trafficking by organellar pH and by proteasome activity. *J Biol Chem* 2004; 279:7971-81.
18. Lamason RL, Mohideen MA, Mest JR, Wong AC, Norton HL, Aros MC, et al. SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* 2005; 310:1782-6.
19. Wolbach SB, Howe PR. Tissue changes following deprivation of fat soluble vitamin A. *J Exp Med* 1925; 43:753.
20. Eckert RL, Broome AM, Ruse M, Robinson N, Ryan D, Lee K. S100 proteins in the epidermis. *J Invest Dermatol* 2004; 123:23-33.